

Physical Aspects of Differential Calculi on Commutative Algebras*

Folkert Müller-Hoissen

Institut für Theoretische Physik
Bunsenstr. 9, D-37073 Göttingen, Germany

Abstract

The central structure in various versions of noncommutative geometry is a differential calculus on an algebra $\mathcal{A}$. This is an analogue of the calculus of differential forms on a manifold. In this short review we collect examples of differential calculi on *commutative algebras* and explain how these are related to relevant structures in physics.

Contents

1	Introduction	2
2	Algebraic differential calculus	2
2.1	The universal differential algebra	3
3	Differential calculus and lattices	3
3.1	The one-dimensional case	3
3.2	Generalization to higher dimensions	5
3.3	Lattice gauge theory	6
3.4	Transformation to q -calculus	7
4	A deformation of the ordinary calculus of differential forms on a manifold	7
4.1	Gauge theory	9
4.2	Generalized differential geometry	9
4.3	Example: Classical limit of bicovariant differential calculus on a quantum group . . .	10
4.4	Relation with ‘proper time’ theories and stochastic calculus on manifolds	11
5	Some aspects of differential calculus on discrete sets	12
5.1	Differential calculus on discrete sets	13
5.2	Reductions of the universal differential algebra, graphs and topologies	14
5.3	Gauge fields	16
5.4	How to recover the usual lattice calculus	17
5.5	‘Symmetric’ differential calculi	17
5.6	The symmetric lattice	18
5.7	Graphy gravity	19
6	Conclusions	19

* Lectures presented at the XXX_q Karpacz Winter School for Theoretical Physics 1994.

1 Introduction

The algebra of differential forms on a manifold $\mathcal{M}$ extends the commutative algebra of C^∞ -functions on a manifold to a differential algebra. Generalizing $C^\infty(\mathcal{M})$ to an arbitrary associative algebra, we may still extend it to a differential algebra keeping the basic properties of the classical exterior derivative (see section 2). This yields a convenient mathematical framework to build physical models even on a ‘noncommutative space’ in close analogy with corresponding ‘classical models’.

Physical motivations to go beyond manifolds and smooth structures in formulating physical dynamics originated in particular from ideas about quantum gravity (see [1], in particular). There have also been suggestions towards a fundamental discreteness of space-time (see [2] for an incomplete list of references). Noncommutative geometry – and in particular algebraic differential calculus – appears to be the appropriate framework for pursuing such ideas.

As an intermediate step towards differential calculus on ‘noncommutative spaces’ we may consider nonstandard differential calculi on commutative spaces. This is what we will concentrate on in these lectures (see also [3]). It will lead us to various familiar structures in physics, but also open routes towards new physical models.

2 Algebraic differential calculus

Let $\mathcal{A}$ be an associative (and not necessarily commutative) algebra (over $\mathbb{R}$ or $\mathbb{C}$). A *differential algebra* $(\Omega(\mathcal{A}), d)$ over $\mathcal{A}$ is a $\mathbb{Z}$ -graded associative algebra

$$\Omega(\mathcal{A}) = \bigoplus_{r=0}^{\infty} \Omega^r(\mathcal{A}) \quad (2.1)$$

where $\Omega^r(\mathcal{A})$ are $\mathcal{A}$ -bimodules and $\Omega^0(\mathcal{A}) = \mathcal{A}$. It is supplied with a linear operator

$$d : \Omega^r(\mathcal{A}) \rightarrow \Omega^{r+1}(\mathcal{A}) \quad (2.2)$$

satisfying $d^2 = 0$ and

$$d(\omega\omega') = (d\omega)\omega' + (-1)^r\omega d\omega' \quad (2.3)$$

where $\omega \in \Omega^r(\mathcal{A})$.

The Grassmann algebra of differential forms on a C^∞ -manifold $\mathcal{M}$ together with the exterior derivative operator d constitutes a differential algebra. Here $\mathcal{A} = C^\infty(\mathcal{M})$, the (commutative) algebra of infinitely often differentiable functions on $\mathcal{M}$. Whereas in this case differentials and functions commute, this need not be so in general for a differential calculus on a commutative algebra. Corresponding examples of *non*-commutative differential calculi on commutative algebras will be discussed in the following sections. In the case of the algebra of functions on a finite set we are actually forced to dispense with commutativity of functions and differentials (see section 5.1).

Given an algebra $\mathcal{A}$, there are many different choices of differential algebras. The problem arises which of them should we choose to work with. What is the significance of different choices? In particular, section 5 will provide us with an interesting answer to this question.

2.1 The universal differential algebra

For any associative algebra $\mathcal{A}$ one can construct the so-called *universal differential algebra*¹ in the following way. We formally associate with every element $a \in \mathcal{A}$ a symbol da and define $\Omega(\mathcal{A})$ to be the linear span over $\mathbb{C}$ of all ‘words’ which can be formed from the symbols a and da ($\forall a \in \mathcal{A}$). Assuming the Leibniz rule $(da)b = d(ab) - a db$ ($\forall a, b \in \mathcal{A}$), each element of $\Omega(\mathcal{A})$ can be written as a sum of monomials of the form $a_0(da_1)\cdots(da_n)$ with $a_i \in \mathcal{A}, i = 1, \dots, n$ and $a_0 \in \mathbb{C}$ or $a_0 \in \mathcal{A}$. It is then sufficient to define the action of d on such monomials:

$$d(a_0 da_1 \cdots da_n) = da_0 da_1 \cdots da_n \quad (2.4)$$

where $da_0 = 0$ if $a_0 \in \mathbb{C}$. As a consequence, we have $d^2 = 0$ and the graded Leibniz rule (2.3). The differential algebra obtained in this way is called *universal differential algebra* of $\mathcal{A}$. If $\mathcal{A}$ has a unit element $\mathbb{1}$ such that $d\mathbb{1} = 0$, there is a representation in terms of tensor products of $\mathcal{A}$ where $da = \mathbb{1} \otimes a - a \otimes \mathbb{1}$ and, more generally,²

$$d(a_0 \otimes a_1 \otimes \cdots \otimes a_n) = \sum_{k=0}^{n+1} (-1)^k a_0 \otimes \cdots \otimes a_{k-1} \otimes \mathbb{1} \otimes a_k \otimes \cdots \otimes a_n. \quad (2.5)$$

Multiplication is defined by concatenation. For example,

$$\begin{aligned} a_0 da_1 da_2 &= (a_0 \otimes a_1 - a_0 a_1 \otimes \mathbb{1}) da_2 = a_0 \otimes a_1 \otimes a_2 - a_0 \otimes a_1 a_2 \otimes \mathbb{1} \\ &\quad - a_0 a_1 \otimes \mathbb{1} \otimes a_2 + a_0 a_1 \otimes a_2 \otimes \mathbb{1}. \end{aligned} \quad (2.6)$$

Any differential algebra over $\mathcal{A}$ can be obtained from the universal differential algebra as a quotient with respect to some differential ideal. The universal differential algebra should therefore be the starting point for a systematic exploration of differential calculi on a given algebra $\mathcal{A}$ (see section 5).

3 Differential calculus and lattices

The simplest deformation of the ordinary differential calculus on $\mathbb{R}^n$ leads to lattice theories. This is the topic of this section which is based on [5, 6].

3.1 The one-dimensional case

A deformation of the ordinary differential calculus on $\mathbb{R}$ is obtained as follows. Let x be the identity function on $\mathbb{R}$ (i.e., $x(r) = r \ \forall r \in \mathbb{R}$) and dx its (formal) differential. We impose the commutation relation

$$[x, dx] = \ell dx \quad (3.1)$$

¹Respectively, *universal differential envelope* of $\mathcal{A}$, see [4] and references given there.

²The argument of d should actually be restricted to an n -form. The tensor product is over $\mathbb{C}$. The space of 1-forms is $\ker m$ where m is the multiplication map $\mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$.

where ℓ is a positive real constant. As a consequence,

$$f(x) dx = dx f(x + \ell) \quad (3.2)$$

for a function of x . Commuting $f(x)$ with dx thus results in a discrete translation of the argument of f . For $\mathcal{A}$ we take the algebra of all functions $\mathbb{R} \rightarrow \mathbb{R}$. They can be regarded as functions of x . Let us assume that dx generates the space $\Omega^1(\mathcal{A})$ of 1-forms as a right- $\mathcal{A}$ -module. Then

$$df = dx \partial_x f \quad (3.3)$$

defines a linear operator $\partial_x : \mathcal{A} \rightarrow \mathcal{A}$. Now

$$\begin{aligned} \ell dx \partial_x f &= [x, dx] \partial_x f = [x, df] \\ &= d(xf - fx) + [f, dx] \quad (\text{using the Leibniz rule for } d) \\ &= f(x) dx - dx f(x) = dx (f(x + \ell) - f(x)) \end{aligned} \quad (3.4)$$

shows that

$$\partial_x f = \frac{1}{\ell} \{f(x + \ell) - f(x)\}. \quad (3.5)$$

dx is also a basis of $\Omega^1(\mathcal{A})$ as a *left*- $\mathcal{A}$ -module, so that

$$df =: \partial_{-x} f dx = \frac{1}{\ell} \{f(x) - f(x - \ell)\} dx. \quad (3.6)$$

Hence d acts as a discrete derivative.³ In particular, it follows that a ‘constant’ h in the sense that $dh = 0$ is a function of x with period ℓ , i.e. $h(x + \ell) = h(x)$. An indefinite integral associated with d is determined by demanding

$$\int df = f + \text{‘constant’}. \quad (3.7)$$

For example,

$$\int dx x = \int dx^2 - \int x dx = x^2 - \ell x - \int dx x \quad (3.8)$$

(modulo a ‘constant’) so that

$$\int dx x = \frac{1}{2} x(x - \ell) + \text{‘constant’}. \quad (3.9)$$

It turns out that every function can be integrated and there is an explicit formula for its indefinite integral [5]. Since ‘constants’ with respect to d are ℓ -periodic functions, a *definite* integral is only defined over intervals of length a multiple of ℓ . In this case one finds [5]

$$\int_{x_0 - m\ell}^{x_0 + n\ell} dx f(x) = \ell \sum_{k=-m}^{n-1} f(x_0 + k\ell) \quad (3.10)$$

($m, n \in \mathbb{N}$). The integral

$$\int_{x_0 - \infty}^{x_0 + \infty} dx f(x) = \sum_{k=-\infty}^{\infty} f(x_0 + k\ell) \quad (3.11)$$

thus picks out the values of f on a *lattice* with ℓ as the lattice spacing.

³In the limit $\ell \rightarrow 0$, the *left*- and *right*-partial derivatives ∂_{-x} and ∂_x tend to the ordinary partial derivative $\partial/\partial x$ (on differentiable functions).

3.2 Generalization to higher dimensions

An obvious generalization of (3.1) to n dimensions is

$$[x^\mu, dx^\nu] = \ell \delta^{\mu\nu} dx^\nu \quad (\text{no summation}) . \quad (3.12)$$

As a consequence,

$$f(x) dx^\mu = dx^\mu f(x + \ell^\mu) \quad (3.13)$$

with the notation $(x + \ell^\mu)^\nu := x^\nu + \delta^{\mu\nu} \ell$. Introducing left- and right-partial derivatives via

$$\sum_{\mu=1}^n \partial_{-\mu} f dx^\mu = df = \sum_{\mu=1}^n dx^\mu \partial_\mu f \quad (3.14)$$

we find

$$\partial_{-\mu} f = \frac{1}{\ell} \{f(x) - f(x - \ell^\mu)\} \quad \partial_\mu f = \frac{1}{\ell} \{f(x + \ell^\mu) - f(x)\} . \quad (3.15)$$

Hence $\partial_{-\mu}$ and ∂_μ are discrete (partial) derivatives.

Again, an indefinite integral is determined by

$$\int df = f + \text{periodic function} \quad (3.16)$$

and a definite integral is only defined over (a union of) cubes with edge length ℓ . In particular, one obtains

$$\int_{x_0^1}^{x_0^1 + \ell} dx^1 \cdots \int_{x_0^n}^{x_0^n + \ell} dx^n f(x^1, \dots, x^n) = \ell^n f(x_0^1, \dots, x_0^n) . \quad (3.17)$$

Using (2.3), a consequence of (3.12) is $dx^\mu dx^\nu = -dx^\nu dx^\mu$. The familiar anticommutation rule for 1-forms thus holds for the differentials dx^μ . It does not hold for arbitrary 1-forms, however. Nevertheless, this allows us to introduce a Hodge operator as follows:

$$\star (dx^{\mu_1} \cdots dx^{\mu_k}) := \frac{1}{(n-k)!} \sum_{\mu_{k+1}, \dots, \mu_n=1}^n \epsilon^{\mu_1 \cdots \mu_k \mu_{k+1} \cdots \mu_n} dx^{\mu_{k+1}} \cdots dx^{\mu_n} \quad (3.18)$$

$$\star (f(x) \omega) := (\star \omega) f(x) . \quad (3.19)$$

Here we have introduced $\epsilon^{\mu_1 \cdots \mu_n}$ which is totally antisymmetric with $\epsilon^{1 \cdots n} = 1$. Indices have been lowered with a Euclidean (or Minkowski) metric with coefficients $\delta_{\mu\nu}$ (or $\eta_{\mu\nu}$) with respect to the coordinate functions x^μ .

We now have all the ingredients for the construction of Lagrangians and actions in terms of differential forms. For a real scalar field ϕ in $n = 4$ dimensions the continuum action can be written as

$$\int -\frac{1}{2} (\star d\phi) d\phi + \frac{1}{2} m^2 (\star \phi) \phi . \quad (3.20)$$

This expression also makes sense for our deformed differential calculus if the integration is understood over a union of cubes with edge length ℓ . It is then easily shown to reproduce the usual lattice action.

The transition from the continuum to a lattice regularizes the corresponding quantum field theory. We see that this is achieved by introducing a noncommutativity between functions and differentials. We do not need ‘noncommuting coordinates’ (i.e. a noncommutative algebra $\mathcal{A}$) to achieve this. Since divergencies appear in *integrated* expressions, noncommutativity between differentials and functions seems to be sufficient.

3.3 Lattice gauge theory

Let $\mathbf{G}$ be a unitary Lie group. A *connection* (or ‘gauge potential’) is a 1-form

$$A = \sum_{\mu=1}^n dx^\mu A_\mu \quad (3.21)$$

which transforms according to the familiar rule

$$A \mapsto G A G^{-1} - dG G^{-1} \quad (3.22)$$

under a gauge transformation with a $\mathbf{G}$ -valued function G . For $\ell = 0$ the gauge potential is Lie algebra valued. For $\ell > 0$, however, it is rather group algebra valued (note that d is a discrete derivative). Then, as a consequence of (3.22), one finds that

$$U := \sum_{\mu} dx^\mu U_\mu := \sum_{\mu} dx^\mu - \ell A \quad (3.23)$$

transforms homogeneously,

$$U \mapsto G U G^{-1} \quad U_\mu(x) \mapsto G(x + \ell^\mu) U_\mu(x) G(x)^{-1} . \quad (3.24)$$

The (gauge covariant) field strength of A is

$$F = dA + A^2 = \frac{1}{\ell^2} \sum_{\mu, \nu} dx^\mu dx^\nu U_\mu(x + \ell^\nu) U_\nu(x) \quad (3.25)$$

and the Yang-Mills action in n dimensions becomes

$$\begin{aligned} S_{YM} &= \int \text{tr}[(\star F^\dagger) F] \\ &= \int d^n x \text{tr} \sum_{\mu, \nu} \ell^{-2n} \left[1 - U_\nu(x)^{-1} U_\mu(x + \ell^\nu)^{-1} U_\nu(x + \ell^\mu) U_\mu(x) \right] \end{aligned} \quad (3.26)$$

where we have made the additional assumption that U_μ has an inverse in $\mathbf{G}$ (which restricts the allowed connections A). When the integral is evaluated over a union of cubes with edge length ℓ , the last expression reproduces the Wilson action of lattice gauge theory.

3.4 Transformation to q -calculus

Let us return to the one-dimensional (deformed) differential calculus of section 3.1 and make a transition to a new (coordinate) function

$$y = q^{x/\ell} \quad (3.27)$$

with $q \in \mathbb{C} \setminus \{0, 1\}$. The basic commutation relation (3.1) is then turned into

$$y \, dy = q \, dy \, y . \quad (3.28)$$

For a function of y we introduce left- and right-partial derivatives with respect to y via

$$\partial_{-y} f \, dy = df = dy \, \partial_y f . \quad (3.29)$$

A simple calculation reveals that these are the q -derivatives

$$\partial_{-y} f = \frac{f(y) - f(q^{-1}y)}{(1 - q^{-1})y} \quad \partial_y f = \frac{f(qy) - f(y)}{(q - 1)y} . \quad (3.30)$$

They span a ‘quantum plane’

$$\partial_{-y} \partial_y = q \, \partial_y \partial_{-y} \quad (3.31)$$

and satisfy q -deformed canonical commutation relations with y (regarded as a multiplication operator),

$$\partial_{-y} y - q^{-1} y \partial_{-y} = 1 \quad \partial_y y - q y \partial_y = 1 . \quad (3.32)$$

Furthermore, the integral introduced in section 3.1 is transformed into the q -integral:

$$\int_{0-\infty}^{0+\infty} dx \dots \mapsto \int_0^\infty dy f(y) = (q - 1) \sum_{k=-\infty}^\infty f(q^k) q^k . \quad (3.33)$$

Here we have to assume that q is not a root of unity.⁴ The q -deformed canonical commutation relations which we obtained above have been considered by various authors. They can be used to formulate a version of q -deformed quantum mechanics (see [5] and the references therein).

4 A deformation of the ordinary calculus of differential forms on a manifold

According to our present knowledge gravity couples to all kind of matter and cannot be shielded away. This universality suggests to build it into the most basic structure needed

⁴The case when $q^N = 1$ for some $N \in \mathbb{N}$ corresponds to taking x modulo $N\ell$. We are then dealing with a periodic lattice.

to formulate physical theories, namely the differential calculus. Along these lines of thought one is led [7] to a differential algebra with

$$[x^i, dx^j] = \tau g^{ij} \quad (4.1)$$

where g^{ij} (which in the above context should be the space-time metric) has to be symmetric as required by the Leibniz rule for d . τ is a 1-form for which we require that

$$[x^i, \tau] = 0 \quad \tau^2 = 0 \quad d\tau = 0. \quad (4.2)$$

Furthermore, we choose $\mathcal{A} = C^\infty(\mathcal{M})$ for a manifold $\mathcal{M}$ and demand that dx^i and τ form a basis of $\Omega^1(\mathcal{A})$ as a right- (or left-) $\mathcal{A}$ -module.⁵ As a consequence, the differential of a function f can be written as

$$df = \tau \hat{\partial}_0 f + dx^i \hat{\partial}_i f \quad (4.3)$$

which defines generalized (right-) partial derivative operators $\hat{\partial}_0, \hat{\partial}_i : \mathcal{A} \rightarrow \mathcal{A}$. Using the Leibniz rule for d , (4.3), the first of the commutation relations (4.2), $[f, dx^i] = [x^i, df]$ (using again the Leibniz rule for d) and (4.1), we find

$$\tau \hat{\partial}_0(fh) + dx^i \hat{\partial}_i(fh) = \tau [(\hat{\partial}_0 f)h + f \hat{\partial}_0 h + g^{ij}(\hat{\partial}_i f)(\hat{\partial}_j h)] + dx^i [(\hat{\partial}_i f)h + f \hat{\partial}_i h]. \quad (4.4)$$

Hence $\hat{\partial}_i$ is a derivation and therefore a vector field (since all derivations of $C^\infty(\mathcal{M})$ are vector fields). (4.3) in particular yields $dx^i = \tau \hat{\partial}_0 x^i + dx^j \hat{\partial}_j x^i$ which implies $\hat{\partial}_0 x^i = 0$ and $\hat{\partial}_j x^i = \delta_j^i$. Taking this into account, we obtain $\hat{\partial}_i = \partial_i$ (the ordinary partial derivative). Writing $\hat{\partial}_0 = \frac{1}{2} g^{ij} \partial_i \partial_j + \delta$, we find that δ must be a derivation and therefore a vector field. But then $\hat{\partial}_0 x^i = 0$ enforces $\delta = 0$. Hence

$$df = \tau \frac{1}{2} g^{ij} \partial_i \partial_j f + dx^i \partial_i f. \quad (4.5)$$

The differential of a function thus involves a *second order* differential operator. Another surprise is that our basic commutation relation (4.1) is invariant under changes of coordinates if g^{ij} are tensor components and τ is a 1-form:

$$\begin{aligned} [x'^i, dx'^j] &= [x'^i, dx^\ell \partial_\ell x'^j + \tau \hat{\partial}_0 x'^j] \\ &= [x'^i, dx^\ell] \partial_\ell x'^j \\ &= [x^\ell, dx'^i] \partial_\ell x'^j \\ &= [x^\ell, dx^k] (\partial_k x'^i) (\partial_\ell x'^j) \\ &= \tau g^{k\ell} (\partial_k x'^i) (\partial_\ell x'^j) \\ &= \tau g'^{k\ell}. \end{aligned} \quad (4.6)$$

The deformed differential calculus is therefore well-defined on a manifold if g and τ are globally defined on $\mathcal{M}$. This opens a way towards some kind of *second order differential geometry*. Corresponding constructions will be presented after a formulation of gauge theory with the deformed differential calculus in the following subsection. What we present here is not a physical theory yet but rather a mathematical framework which we expect to be useful for the formulation and investigation of physical theories.

⁵This structure is chosen as to cover some examples to which we will turn in subsections. These examples must not necessarily be put into the physical context outlined in the beginning of this section.

4.1 Gauge theory

Let ψ be an element of $\mathcal{A}^n$ which transforms as $\psi \mapsto \psi' = G\psi$ under a representation of a Lie group. For local transformations we can construct a covariant derivative in the usual way,

$$D\psi = d\psi + A\psi. \quad (4.7)$$

This is indeed covariant if the 1-form A transforms according to the familiar rule

$$A' = G A G^{-1} - dG G^{-1}. \quad (4.8)$$

A can then be written in a unique way as

$$A = \tau \frac{1}{2} A_\tau + dx^i A_i. \quad (4.9)$$

Inserting this expression in (4.8), we find that A_i behaves as an ordinary gauge potential and

$$A_\tau = g^{ij} (\partial_i A_j - A_i A_j) + M \quad (4.10)$$

where M is an arbitrary tensorial part ($M' = G M G^{-1}$). In order to be able to read off gauge-covariant components from gauge-covariant differential forms, we need the covariantized differentials $Dx^i := dx^i - \tau A^i$. They transform as $D'x^i = G Dx^i G^{-1}$. The covariant derivative of ψ can now be written as

$$D\psi = \tau \frac{1}{2} (g^{ij} D_i D_j + M) \psi + Dx^i D_i \psi \quad (4.11)$$

where D_i denotes the ordinary covariant derivative (using A_i). The field strength of A is

$$F = dA + A^2 = \tau \frac{1}{2} (dx^i D^j F_{ji} - DM) + \frac{1}{2} Dx^i Dx^j F_{ij} \quad (4.12)$$

which involves the Yang-Mills operator (when g^{ij} is identified with the space-time metric). F_{ij} is the (ordinary) field strength of A_i .

4.2 Generalized differential geometry

For a vector field Y^i we introduce a (right-) covariant derivative

$$DY^i := dY^i + Y^j {}_j\Gamma^i. \quad (4.13)$$

This is indeed *right*-covariant iff the generalized connection ${}_j\Gamma^i$ is given by

$${}_j\Gamma^i = \tau \frac{1}{2} \left[g^{k\ell} (\partial_k \Gamma^i_{j\ell} + \Gamma^i_{mk} \Gamma^m_{j\ell}) + M^i_j \right] + dx^k \Gamma^i_{jk} \quad (4.14)$$

where Γ^i_{jk} are the components of an ordinary linear connection on $\mathcal{M}$ and M^i_j is a tensor.

The coordinate differentials dx^i do not transform covariantly, since

$$dx'^k = \tau \frac{1}{2} g^{ij} \partial_i \partial_j x'^k + dx^\ell \partial_\ell x'^k \quad (4.15)$$

as a consequence of (4.5). But the 1-forms

$$Dx^k := dx^k + \tau \frac{1}{2} \Gamma^k_{ij} g^{ij} \quad (4.16)$$

are *right*-covariant and (4.5) can now be rewritten as

$$df = \tau \frac{1}{2} g^{ij} \nabla_i \nabla_j f + Dx^i \partial_i f \quad (4.17)$$

where ∇_i denotes the ordinary covariant derivative (with $\Gamma^i_{k\ell}$). Also the covariant exterior derivative of Y^i can now be written in an explicitly right-covariant form,

$$DY^i = \tau \frac{1}{2} (g^{k\ell} \nabla_k \nabla_\ell Y^i + M^i_j Y^j) + Dx^j \nabla_j Y^i. \quad (4.18)$$

The product of differentials Dx^k is neither antisymmetric nor covariant, but

$$Dx^k \wedge Dx^\ell := Dx^k Dx^\ell - \tau dx^j g^{i\ell} \Gamma^k_{ij} \quad (4.19)$$

is antisymmetric and right-covariant. Then

$$D^2 x^k = \Theta^k - \tau \frac{1}{2} dx^\ell (R^k_\ell - M^k_\ell - \nabla^j Q^k_{j\ell} + Q^{kij} Q_{ij\ell}) \quad (4.20)$$

with the torsion 2-form

$$\Theta^k := \frac{1}{2} Dx^i \wedge Dx^j Q^k_{ij} \quad (4.21)$$

where Q^k_{ij} is the torsion of Γ^k_{ij} . (4.20) may be regarded as a generalized torsion. Its vanishing implies the vanishing of Q^k_{ij} and an equation of the form of Einstein's equations. It suggests to relate M^i_j to the energy-momentum tensor of matter. The latter is then 'geometrized' in this formalism as part of the generalized connection.

We have seen in particular that the (covariant) exterior derivative of a field contains in its τ -part the corresponding part of the field equation to which it is usually subjected in physical models. We refer to [7] for an ample discussion.

4.3 Example: Classical limit of bicovariant differential calculus on a quantum group

The Hopf algebra structure of a quantum group can be used to narrow down the large number of possible differential calculi on such an algebra and leads to the concept of 'bicovariant differential calculus' [8]. One is mainly interested in such calculi where the dimension of the space of 1-forms (as a left or right $\mathcal{A}$ -module) coincides with the number of generators of the quantum group.

The quantum group $GL_q(2)$ is an algebra $\mathcal{A}$ generated by elements x^μ , $\mu = 1, \dots, 4$, which satisfy certain commutation relations. All possible bicovariant differential calculi on $GL_q(2)$ were found in [9] (see also [10]). They form a 1-parameter family (with a complex parameter s). In the classical limit $q \rightarrow 1$ one obtains [11]⁶

$$[x^\mu, dx^\nu] = \tau \gamma^{\mu\nu} \quad (4.22)$$

with

$$\gamma^{\mu\nu} = -(x^1 x^4 - x^2 x^3)^{-1} x^\mu x^\nu + 4(\delta_1^{(\mu} \delta_4^{\nu)} - \delta_2^{(\mu} \delta_3^{\nu)}) \quad (4.23)$$

$$\tau = s(x^4 dx^1 - x^3 dx^2 - x^2 dx^3 + x^1 dx^4) =: \tau_\nu dx^\nu. \quad (4.24)$$

These objects satisfy $\gamma^{\mu\nu} \tau_\nu = 0$ which defines a Galilei structure⁷.

Passing from differential calculus on $GL_q(2)$ to differential calculus on $SL_q(2, \mathbb{R})$ one has to fix the parameter s . From the two resulting bicovariant differential calculi only one has a reasonable classical limit. After elimination of x^4 with the help of the determinant constraint in a coordinate patch where $x^1 \neq 0$, we obtain [10]

$$[x^i, dx^j] = \tau g^{ij} \quad (i, j = 1, 2, 3) \quad (4.25)$$

where g^{ij} is the bi-invariant metric on $SL(2, \mathbb{R})$. The 1-form τ cannot be expressed as $\sum_i f_i(x^j) dx^i$ or $\sum_i dx^i f_i(x^j)$. The 1-forms dx^i and τ thus form a basis of $\Omega^1(\mathcal{A})$. We refer to [10, 11, 12] for further details and to [13] for a generalization of the above example.

4.4 Relation with ‘proper time’ theories and stochastic calculus on manifolds

When $\tau = \gamma dt$ with a constant γ , we may consider (smooth) functions $f(x^i, t)$ depending also on the parameter t . (4.5) then has to be replaced by

$$df = dt \left(\partial_t + \frac{\gamma}{2} g^{ij} \partial_i \partial_j \right) f + dx^i \partial_i f. \quad (4.26)$$

1. Let $\gamma = -i\hbar$. The requirement that the dt -part vanishes leads to the Schrödinger equation

$$(i\hbar \partial_t + \frac{\hbar^2}{2} g^{ij} \partial_i \partial_j) f = 0. \quad (4.27)$$

If g^{ij} is a space-time metric, this is the five-dimensional Schrödinger equation of ‘proper time’ quantum theory (à la Fock, Stueckelberg, Nambu, Feynman, ..., see [14] for a recent review).

⁶The limit was obtained by regarding the parameter $s = A_4^1$ in the notation of [9, 10] as independent of q . In this parametrization there are two calculi for a given value of s . Here we refer to the one for which $A_1^1 = 1 + s$ (in the notation of the cited papers). After a q -dependent transformation to another parameter t (as in [12]) we may as well regard this new parameter t as q -independent. Then the limit $q \rightarrow 1$ yields in general a different result. This is in fact the case with the parameter t chosen in [12]. The statement made there about how the limit was taken is incorrect.

⁷This is a generalization of the Newtonian space-time in which case $\gamma^{\mu\nu}$ is the space-metric and $\tau = dt$ where t is the absolute time.

The noncommutative differential calculus may be viewed as a basic structure underlying such proper time theories (see [7] for an extensive discussion).

2. The formula (4.26) – with a positive definite metric γg^{ij} – is well-known in the theory of stochastic processes (Itô calculus) and suggests that the noncommutative differential calculus provides us with a convenient framework to deal with stochastic processes on manifolds (cf [15] where $\gamma = \hbar$ and g is the inverse mass matrix). There is indeed a translation [11] to the (Itô) calculus of (commutative) stochastic differentials where one has the following rules for products of stochastic differentials:

$$dt \bullet dt = 0, \quad dt \bullet dx_t^i = 0, \quad dx_t^i \bullet dx_t^j = \gamma g^{ij} dt. \quad (4.28)$$

The coordinates x^i have been replaced by stochastic processes x_t^i and g^{ij} is a function of the latter. In the stochastic calculus one has a deformed Leibniz rule:

$$d(f_t h_t) = (df_t) h_t + f_t dh_t + (df_t) \bullet dh_t. \quad (4.29)$$

But it is possible to define a new (noncommutative) product $*$ between functions and differentials such that

$$d(f_t h_t) = (df_t) * h_t + f_t * dh_t. \quad (4.30)$$

(This product is related to the Moyal product.) In terms of the corresponding commutator the relations (4.28) are translated into

$$[t, dt]_* = 0, \quad [t, dx_t^i]_* = [x_t^i, dt]_* = 0, \quad [x_t^i, dx_t^j]_* = \gamma g^{ij} dt. \quad (4.31)$$

In contrast to the Itô calculus, the noncommutative differential calculus admits an extension to forms of higher grade. Do they have a stochastic interpretation? We refer to [11, 13] for further details. The formalism should also be of interest in the context of stochastic quantization.

5 Some aspects of differential calculus on discrete sets

When one formulates differential calculus on (the commutative algebra of functions on) a finite or, more generally, discrete set, functions and differentials turn out to be necessarily *noncommutative*. The simplest example is provided by Connes' 2-point model which received a lot of interest as an ingredient of particle physics models [16]. Another example, a lattice differential calculus, was already discussed in section 3. These two examples motivated a general study of differential calculi on arbitrary discrete sets. This has been carried out in some recent papers [17, 18] (see also [19] for the case of discrete groups). We believe that this kind of mathematics is very convenient for discrete modelling and in particular for developing ideas of discrete space-time (cf [20]).

5.1 Differential calculus on discrete sets

Let $\mathcal{M}$ be a discrete⁸ set of elements $i, j, \dots$ and $\mathcal{A}$ the algebra of $\mathbb{C}$ -valued functions on it. With each element $i \in \mathcal{M}$ we associate a function $e_i \in \mathcal{A}$ via

$$e_i(j) = \delta_{ij} . \quad (5.1)$$

Then $e_i e_j = \delta_{ij} e_j$ and $\sum_i e_i = \mathbb{I}$ where $\mathbb{I}(i) = 1 \ \forall i \in \mathcal{M}$. Acting with d on these relations and using the Leibniz rule yields

$$e_i de_j = -de_i e_j + \delta_{ij} de_j \quad \sum_i de_i = 0 \quad (5.2)$$

where the first formula expresses the *noncommutativity* of the differential calculus. It turns out to be convenient to introduce the following 1-forms,

$$e_{ij} := e_i de_j \quad (i \neq j) \quad (5.3)$$

(we set $e_{ii} = 0$). From these 1-forms we build the $(r-1)$ -forms

$$e_{i_1 \dots i_r} := e_{i_1 i_2} e_{i_2 i_3} \dots e_{i_{r-1} i_r} . \quad (5.4)$$

Now one has the following formulae,

$$de_i = \sum_k (e_{ki} - e_{ik}) \quad df = \sum_{i,j} e_{ij} [f(j) - f(i)] \quad (5.5)$$

$$de_{ij} = \sum_k (e_{kij} - e_{ikj} + e_{ijk}) . \quad (5.6)$$

If no further relations are imposed, we are dealing with the universal differential calculus. In this case the $e_{i_1 \dots i_r}$ constitute a basis over $\mathbb{C}$ of $\Omega^{r-1}(\mathcal{A})$ for $r > 1$.

An *inner product* on $\Omega(\mathcal{A})$ should have the properties

$$(\psi_r, \phi_s) = 0 \quad \text{if } r \neq s \quad (5.7)$$

$$(\psi, c\phi + \chi) = c(\psi, \phi) + (\psi, \chi) \quad \forall c \in \mathbb{C} \quad (5.8)$$

$$\overline{(\psi, \phi)} = (\phi, \psi) \quad (5.9)$$

(where $\psi_r \in \Omega^r(\mathcal{A})$ and the bar denotes complex conjugation). An inner product is then determined by the values of $(e_{i_1 \dots i_r}, e_{j_1 \dots j_r})$. We will require that this has the structure

$$(e_{i_1 \dots i_r}, e_{j_1 \dots j_r}) = \delta_{i_1 j_1} g_{i_1 \dots i_r j_1 \dots j_r} \delta_{i_r j_r} \quad (5.10)$$

with constants $g_{i_1 \dots i_r j_1 \dots j_r}$. For the universal $\Omega(\mathcal{A})$, the latter may be taken proportional to $\delta_{i_1 j_1} \dots \delta_{i_r j_r} \prod_{s=1}^{r-1} (1 - \delta_{i_s i_{s+1}})$ (the last factor takes care of the fact that $e_{ii} = 0$). Given an inner product, we can construct an adjoint $d^\dagger$ of d , a Laplace-Beltrami operator etc..

An *involution* on $\Omega(\mathcal{A})$ should have the properties

$$(\psi \phi)^* = \phi^* \psi^* \quad (d\psi_r)^* = (-1)^r d(\psi_r^*) . \quad (5.11)$$

If $*$ restricted to $\Omega^0(\mathcal{A}) = \mathcal{A}$ is complex conjugation, then

$$e_{kl}^* = -e_{lk} . \quad (5.12)$$

⁸In the case of *infinite* sets the following calculus is ‘formal’ and more work has to be done to put it on a rigorous footing.

5.2 Reductions of the universal differential algebra, graphs and topologies

Smaller differential algebras are obtained from the universal one by setting some of the e_{ij} to zero ('reduction'). Let us associate a *graph* with a differential calculus on $\mathcal{M}$ in the following way. The vertices correspond to the elements of $\mathcal{M}$ and there is an arrow from i to j iff $e_{ij} \neq 0$.

Example 1. The universal differential calculus on a 3-point set is represented by the graph




Fig.1

The graph associated with the universal differential algebra on a set with three elements.

Example 2. Let us consider the differential calculus on the 3-point space associated with the following graph.




Fig.2

The graph associated with a reduction of the universal differential algebra on a set of three elements.

The nonvanishing basic 1-forms are thus e_{01}, e_{12}, e_{20} . The only basic 2-forms we can construct from these are e_{012}, e_{120} and e_{201} according to (5.4). But $e_{10} = e_{02} = e_{21} = 0$ implies

$$0 = de_{10} = \sum_{k=1}^3 (e_{k10} - e_{1k0} + e_{10k}) = -e_{120} \quad (5.13)$$

and also $e_{012} = e_{201} = 0$ so that there are no 2-forms. Hence we can assign the *dimension* 1 to the 3-point set. Let us associate new vertices with the nonvanishing basic 1-forms and draw an extended graph (Fig.3).




Fig.3

The extension of the graph in Fig.2 drawn in two different ways. The second version suggests an interpretation as an (oriented) Hasse diagram.

Following Sorkin [20], the second graph in Fig.3 can be interpreted as a *Hasse diagram* which

determines a topology on the 3-point space in the following way. A vertex together with all lower lying vertices which are connected to it forms an open set. In the present case, $\{01\}, \{12\}, \{20\}, \{0, 01, 20\}, \{1, 01, 12\}, \{2, 12, 20\}$ are the open sets (besides the empty and the whole set). This is an approximation to the topology of S^1 . It consists of a chain of three open sets covering S^1 which already displays the global topology of S^1 . In particular, the fundamental group π_1 is the same as for S^1 .

Example 3. On a 4-point set let us consider the differential calculus associated with the following graph.




Fig.4

A graph representing a differential calculus on a set of four elements.

The nonvanishing basic 1-forms are $e_{01}, e_{12}, e_{03}, e_{32}$. From these we can only construct the basic 2-forms e_{012} and e_{032} . No forms of higher grade are present. Furthermore, (5.6) together with $e_{02} = 0$ implies $e_{032} = -e_{012}$ and there remains only one independent basic 2-form (which we will take to be e_{012}). The differential calculus thus assigns 2 dimensions to the 4-point set. Again, we extend the graph by adding new vertices corresponding to the nonvanishing basic 1- and 2-forms (Fig.5).




Fig.5

The extended graph and (oriented) Hasse diagram derived from the graph in Fig.4.

The arrows are determined by the equations

$$de_{01} = de_{12} = -de_{03} = -de_{32} = e_{012} . \quad (5.14)$$

For example, e_{012} appears in the expression for de_{01} . So we connect the corresponding vertices. The orientation of the arrow is determined by the sign in front of e_{012} . For the topology the orientation is irrelevant, however. The latter can be visualized as follows (Fig.6).




Fig.6

The topology on the 4-point set determined by the Hasse diagram in Fig.5. The four disks and their intersections represent 2-dimensional open sets.

Further examples can be found in [18].

Remark. In a recent work [21] the authors start with a topology (e.g., given by imprecise space-time measurements), construct the Hasse diagram, assign incidence numbers ± 1 to its edges, construct a boundary operator etc.. Using the results outlined above, one recovers a noncommutative differential calculus (on a finite set) behind all this. $\square$

5.3 Gauge fields

As an example of field theory on a discrete set we consider (pure) gauge theory. The structures introduced in the following are defined for any choice of a differential calculus on a discrete set $\mathcal{M}$. A *connection 1-form* is an element $A = \sum_{i,j} e_{ij} A_{ij} \in \Omega^1(\mathcal{A}) \otimes_{\mathcal{A}} M_n(\mathcal{A})$ (where $M_n(\mathcal{A})$ is the space of $n \times n$ matrices with entries in $\mathcal{A}$) with the familiar transformation rule⁹

$$A \mapsto G A G^{-1} - dG G^{-1} \quad (5.15)$$

where $G = \sum_i G(i) e_i$ is an element of a local gauge group, a subgroup of $GL(n, \mathcal{A})$. Associated with A is the *transport operator*

$$U = \sum_{i,j} e_{ij} U_{ij}, \quad U_{ij} := \mathbf{1} + A_{ij} \quad (5.16)$$

which transforms as follows,

$$U \mapsto G U G^{-1} \quad U_{ij} \mapsto G(i) U_{ij} G(j)^{-1}. \quad (5.17)$$

The curvature (or field strength) of A is

$$F := dA + A^2 = \sum_{i,j,k} e_{ijk} (U_{ij} U_{jk} - U_{ik}). \quad (5.18)$$

It transforms in the familiar way, $F \mapsto G F G^{-1}$. In order to generalize an inner product (with the property (5.10)) to matrix valued forms, we require that¹⁰

$$(\phi, \psi) = \sum \phi_{i_1 \dots i_r}^\dagger (e_{i_1 \dots i_r}, e_{j_1 \dots j_s}) \psi_{j_1 \dots j_s}. \quad (5.19)$$

The *Yang-Mills action*

$$S_{YM} := \text{tr} (F, F) \quad (5.20)$$

is then gauge-invariant if $G^\dagger = G^{-1}$. Covariant derivatives of fields on $\mathcal{M}$ can be introduced in the usual way [18].

⁹Note that A cannot be Lie algebra valued since dG is a discrete derivative. See also section 3.3.

¹⁰Here $\phi_{i_1 \dots i_r}$ is a matrix with entries in $\mathbb{C}$ and $\phi_{i_1 \dots i_r}^\dagger$ denotes the hermitian conjugate matrix.

5.4 How to recover the usual lattice calculus

Choose $\mathcal{M} = \mathbb{Z}^n = \{a = (a^\mu) \mid \mu = 1, \dots, n, a^\mu \in \mathbb{Z}\}$ and consider the following reduction of the universal differential calculus on $\mathcal{M}$:

$$e_{ab} \neq 0 \quad \Leftrightarrow \quad b = a + \hat{\mu} \text{ for some } \mu \quad (5.21)$$

where $\hat{\mu} := (\delta_\mu^\nu) \in \mathcal{M}$. The corresponding graph is an oriented lattice in n dimensions (which locally looks like the graph in Fig.4). In terms of the n functions

$$x^\mu := \ell \sum_a a^\mu e_a \quad (5.22)$$

(where ℓ is a positive constant) one finds the relation¹¹

$$dx^\mu f(x) = f(x + \ell \hat{\mu}) dx^\mu \quad (5.23)$$

which shows that we are dealing with the differential calculus of section 3 which led us to lattice (gauge) theories.¹² ℓ plays the role of the lattice spacing.

5.5 ‘Symmetric’ differential calculi

The involution introduced in section 5.1 is defined in a natural way on the universal differential algebra. It is not consistent, in general, with reductions of it since (5.12) requires that with $e_{ij} = 0$ (for some i, j) we must also have $e_{ji} = 0$. A differential calculus on $\mathcal{M}$ with this property (and also the associated graph) is called *symmetric*. For such a symmetric calculus the hermitian conjugation of complex matrices extends to matrix valued differential forms via

$$\phi^\dagger = \sum_{i_1 \dots i_r} (\phi_{i_1 \dots i_r} e_{i_1 \dots i_r})^\dagger = \sum_{i_1 \dots i_r} \phi_{i_1 \dots i_r}^\dagger e_{i_1 \dots i_r}^* . \quad (5.24)$$

The condition $A^\dagger = -A$ for a connection 1-form is then equivalent to $U_{ij}^\dagger = U_{ji}$.

Example. For $\mathcal{M} = \mathbb{Z}_2 = \{0, 1\}$ with the universal differential algebra one finds

$$F = e_{010} (U_{01} U_{10} - \mathbf{1}) + e_{101} (U_{10} U_{01} - \mathbf{1}) = e_{010} (U_{10}^\dagger U_{10} - \mathbf{1}) + e_{101} (U_{10} U_{10}^\dagger - \mathbf{1}) \quad (5.25)$$

using the above condition for A . Then $S_{YM} = 2 \text{tr} (U_{10}^\dagger U_{10} - \mathbf{1})^2$ which has the form of a Higgs potential (cf [16]).

Remark. Using the naive notion of dimension introduced in section 5.2, symmetric graphs are ∞ -dimensional. This is so because from e_{ij} and e_{ji} one can construct forms of arbitrarily high grade: $e_{ijijij\dots}$. Perhaps one should ignore such forms in determining the dimension. Then, for example, the graph shown in Fig.7 is quasi 1-dimensional.

¹¹Every $f \in \mathcal{A}$ can be regarded as a function of x^μ .

¹²In section 3 we considered the last relation (with $\ell \mapsto -\ell$) on the algebra of functions on $\mathbb{R}^n$ and found that it actually lives on a lattice. In contrast, in the present section we started with the algebra of functions on the lattice (i.e., $\mathbb{Z}^n$).




Fig.7

A quasi-1-dimensional graph. The edges represent pairs of antiparallel arrows.

5.6 The symmetric lattice

A particular example of a symmetric differential calculus is defined as follows. We take $\mathcal{M} = \mathbb{Z}^n$ and define a differential calculus by

$$e_{ab} \neq 0 \quad \Leftrightarrow \quad b = a + \hat{\mu} \quad \text{or} \quad b = a - \hat{\mu} \quad \text{for some } \mu \quad (5.26)$$

where $\hat{\mu} = (\delta_\mu^\nu)$ and $\mu = 1, \dots, n$. This is represented by a hypercubic lattice graph where both arrows are present between connected vertices. We therefore call it the *symmetric lattice*. It has no distinguished directions in contrast to the oriented lattice graph (cf section 5.4). Furthermore, it is a lattice generalization of Connes' 2-point space. A technical advantage over the oriented lattice is that the symmetric lattice is compatible with the natural involution defined in section 5.1 (see also section 5.5).

As in the case of the oriented lattice we introduce

$$x^\mu := \ell \sum_a a^\mu e_a . \quad (5.27)$$

Then

$$dx^\mu = \ell \sum_a (e_{a, a+\hat{\mu}} - e_{a, a-\hat{\mu}}) . \quad (5.28)$$

Together with

$$\tau^\mu := \beta \sum_a (e_{a, a+\hat{\mu}} + e_{a, a-\hat{\mu}}) \quad (5.29)$$

(where $\beta \neq 0$ is a constant) the dx^μ constitute a basis of $\Omega^1(\mathcal{A})$ as a left (or right) $\mathcal{A}$ -module. For $f \in \mathcal{A}$ (which can be regarded as a function of x^μ) one finds the relation

$$df = \sum_\mu (\bar{\partial}_\mu f dx^\mu + \frac{\kappa}{2} \Delta_\mu f \tau^\mu) . \quad (5.30)$$

Here $\kappa := \ell^2/\beta$ and we have introduced the operators

$$\partial_{\pm\mu} f := \pm \frac{1}{\ell} (f(x \pm \ell \hat{\mu}) - f(x)) , \quad \bar{\partial}_\mu f := \frac{1}{2} (\partial_{+\mu} f + \partial_{-\mu} f) , \quad \Delta_\mu f := \partial_{+\mu} \partial_{-\mu} f . \quad (5.31)$$

For the commutation relations between functions and 1-forms we find

$$[dx^\mu, f(x)] = \frac{\kappa\beta}{2} \Delta_\mu f(x) dx^\mu + \kappa \bar{\partial}_\mu f(x) \tau^\mu \quad (5.32)$$

$$[\tau^\mu, f(x)] = \beta \bar{\partial}_\mu f(x) dx^\mu + \frac{\kappa\beta}{2} \Delta_\mu f(x) \tau^\mu . \quad (5.33)$$

Especially (5.30) should remind us of a similar formula in section 4.

5.7 Graphy gravity

In this section we anticipate a bit from a forthcoming paper [22]. A *vielbein field* on $\mathcal{M}$ is a 1-form

$$\vartheta = \sum_{i,j} e_{ij} \vartheta_{ij} \quad (5.34)$$

where $\vartheta_{ij} \in \mathbb{R}^n$. It is subject to local $SO(1, n-1)$ transformations,

$$\vartheta \mapsto G \vartheta \quad (5.35)$$

which implies $\vartheta_{ij} \mapsto G(i) \vartheta_{ij}$. We should therefore regard ϑ_{ij} as sitting at the point i . Using $\eta := \text{diag}(-1, 1, \dots, 1)$ we define

$$(\vartheta_{ij}^\dagger)_\mu := \eta_{\mu\nu} \vartheta_{ij}^\nu \quad (\mu = 1, \dots, n) . \quad (5.36)$$

Restricting to symmetric differential calculi (in the sense of section 5.5), we have a natural involution on $\Omega(\mathcal{A})$. With

$$\vartheta^\dagger := \sum_{i,j} e_{ij}^* \vartheta_{ij}^\dagger = - \sum_{i,j} e_{ij} \vartheta_{ji}^\dagger \quad (5.37)$$

we find that $\vartheta \vartheta^\dagger \mapsto G \vartheta \vartheta^\dagger G^{-1}$ under a gauge transformation. If F is the curvature of an $SO(1, n-1)$ connection on $\mathcal{M}$, we can now construct an analogue of the Einstein-Cartan action, namely

$$S_{EC} = \text{tr}(\vartheta \vartheta^\dagger, F) . \quad (5.38)$$

This is indeed gauge-invariant. But what has it to do with gravity?

6 Conclusions

In section 3 we have demonstrated that a simple deformation of the algebra of differential forms on $\mathbb{R}^n$ leads us from continuum to lattice theories. In particular, the usual action for lattice gauge theory is obtained from the continuum Yang-Mills action via this deformation. The ‘lattice differential calculus’ of section 3 can also be regarded as a differential calculus on the set of lattice points (i.e., $\mathbb{Z}^n$). In the language of section 5 it corresponds to a hypercubic lattice graph and we have seen how everything can be generalized to those graphs representing differential algebras on a discrete set (which can be obtained as ‘reductions’ of the universal differential algebra). We have shown how the choice of a differential algebra on a discrete set determines a topology and assigns a dimension to it (which is actually a local notion since it may vary from subgraph to subgraph).

In this sense, a space like $\mathcal{M} \times \mathbb{Z}_2$ (with a manifold $\mathcal{M}$ and the universal differential calculus on $\mathbb{Z}_2$) can be regarded as an approximation of $\mathcal{M} \times S^2$. The origin of a Higgs field in the Connes & Lott models [16] then seems to be quite the same as the origin of Higgs fields in models of dimensional reduction of gauge fields (see [23], in particular). This correspondence still has to be made more precise.

Beyond what we discussed so far, a further aspect seems to be very promising. One can imagine that a specific differential calculus will be determined *dynamically* as a reduction of the universal differential calculus (on a given set). This leads to a framework in which topology change and fluctuations of dimension will naturally occur, features which one should expect in quantum gravity.

Moreover, the differential calculus discussed in section 4 exhibited surprising relations between bicovariant differential calculus on quantum groups, the ‘proper time formalism’ of quantum theory and stochastics. It also provided us with an example of how dynamics can be encoded in the differential calculus and then induced on various fields.

Acknowledgments. I am grateful to Jerzy Lukierski for the kind invitation and to all the organizers for the pleasant atmosphere at the Karpacz winter school. Furthermore, I have to thank Aristophanes Dimakis for a very fruitful and exciting collaboration.

References

- [1] C.J. Isham, *An introduction to general topology and quantum topology*, in: Physics, Geometry and Topology, ed. H.C. Lee (Plenum Press, New York, 1990).
- [2] D. Finkelstein, *Space-time code*, Phys. Rev. **184** (1969) 1261; R.P. Feynman, *Simulating physics with computers*, Int. J. Theor. Phys. **21** (1982) 467; G. Feinberg, R. Friedberg, T.D. Lee and H.C. Ren, *Lattice gravity near the continuum*, Nucl. Phys. B **245** (1984) 343; T.D. Lee, *Discrete mechanics*, in: How far are we from the gauge forces, ed. A. Zichichi (Plenum Press, New York, 1985) p.15; L. Bombelli, J. Lee, D. Meyer and R.D. Sorkin, *Space-time as a causal set*, Phys. Rev. Lett. **59** (1987) 521; G. ’t Hooft, *Quantization of discrete deterministic theories by Hilbert space extension*, Nucl. Phys. B **342** (1990) 471; D. Finkelstein and J.M. Gibbs, *Quantum relativity*, Int. J. Theor. Phys. **32** (1993) 1801.
- [3] F. Müller-Hoissen, *Some applications of noncommutative differential calculus in physics*, habilitation thesis, Göttingen (1992).
- [4] R. Coquereaux, *Noncommutative geometry and theoretical physics*, J. Geom. Phys. **6** (1989) 425.
- [5] A. Dimakis and F. Müller-Hoissen, *Quantum mechanics on a lattice and q -deformations*, Phys. Lett. **295B** (1992) 242.
- [6] A. Dimakis, F. Müller-Hoissen and T. Striker, *From continuum to lattice theory via deformation of the differential calculus*, Phys. Lett. **300B** (1993) 141; *Noncommutative differential calculus and lattice gauge theory*, J. Phys. A **26** (1993) 1927.
- [7] A. Dimakis and F. Müller-Hoissen, *A noncommutative differential calculus and its relation to gauge theory and gravitation*, Int. J. Mod. Phys. A (Proc. Suppl.) **3A** (1993) 474; *Noncommutative differential calculus, gauge theory and gravitation*, Preprint Göttingen GOE-TP 33/92, (1992).
- [8] S.L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups)*, Commun. Math. Phys. **122** (1989) 125.

- [9] F. Müller-Hoissen, *Differential calculi on the quantum group $GL_{p,q}(2)$* , J. Phys. A (Math. Gen.) **25** (1992) 1703.
- [10] F. Müller-Hoissen and C. Reuten, *Bicovariant differential calculus on $GL_{p,q}(2)$ and quantum subgroups*, J. Phys. A (Math. Gen.) **26** (1993) 2955.
- [11] A. Dimakis and F. Müller-Hoissen, *Stochastic differential calculus, the Moyal $\ast$ -product, and noncommutative geometry*, Lett. Math. Phys. **28** (1993) 123.
- [12] F. Müller-Hoissen, *Differential calculi on quantum (sub-) groups and their classical limit*, hep-th/9401152, to appear in the proceedings of the workshop on ‘Generalized Symmetries in Physics’, ASI Clausthal, July 1993.
- [13] A. Dimakis and F. Müller-Hoissen, *Noncommutative Differential Calculus: Quantum Groups, Stochastic Processes, and the Antibracket*, preprint GOET-TP 55/93, hep-th/9401151.
- [14] J. R. Fanchi, *Review of invariant time formulations of relativistic quantum theories*, Found. Phys. **23** (1993) 487.
- [15] E. Nelson, *Quantum Fluctuations* (Princeton University Press, Princeton, 1985).
- [16] A. Connes and J. Lott, *Particle models and noncommutative geometry*, Nucl. Phys. B (Proc. Suppl.) **18** (1990) 29; *The metric aspect of noncommutative geometry*, in: New Symmetry Principles in Quantum Field Theory, eds. J. Fröhlich et al. (Plenum Press, New York, 1992) 53.
- [17] A. Dimakis and F. Müller-Hoissen, *Differential calculus and gauge theory on finite sets*, J. Phys. A: Math. Gen. **27** (1994) 3159. *Differential calculus and discrete structures*, hep-th/9401150, to appear in the proceedings of the workshop on ‘Generalized Symmetries in Physics’, ASI Clausthal, July 1993.
- [18] A. Dimakis and F. Müller-Hoissen, *Discrete differential calculus, graphs, topologies and gauge theory*, preprint GOET-TP 1/94, hep-th/9404112.
- [19] A. Sitarz, *Noncommutative geometry and gauge theory on discrete groups*, preprint TPJU-7/92.
- [20] R. D. Sorkin, *Finitary substitute for continuous topology*, Int. J. Theor. Phys. **30** (1991) 923.
- [21] A. P. Balachandran, G. Bimonte, E. Ercolessi and P. Teotonio-Sobrinho, *Finite approximations to quantum physics: quantum points and their bundles*, Syracuse preprint SU-4240-550, October 1993.
- [22] A. Dimakis and F. Müller-Hoissen, *Noncommutative geometry and gravitation*, in preparation.
- [23] N.S. Manton, *A new six-dimensional approach to the Weinberg-Salam model*, Nucl. Phys. B **158** (1979) 141.